FRUIT FLY CONTROL IN YOUR GARDEN IS A YEAR-ROUND ACTIVITY

START YOUR MONITORING PROGRAM 6-8 WEEKS BEFORE RIPENING BY INSTALLING TRAPS; CONTINUE MONITORING UNTIL 2-3 WEEKS AFTER PICKING.



FRUIT	AUG				SEP				T	ост				NOV				Γ	DEC				JAN			FEB				MAR				APR				MAY				JUN					JUL			
Loquat		0					0			9	9	0	•	0	9		0	•		Ü												П				Г						Г	Г				Γ		Γ	
Apricot					0			0		9		0	•	0	0	•	0	9) (•	0									Г							Г					Γ				Γ		Γ	
Valencia orange	•			•	0	•	9	0		D	•	0	0	0	0	0	0	6) ()	0	0		•																									•	•
Peach		0								0	9	0	•		0	•	0	0)	9	0	•	•	0	0	0	0	•	0	•															3 - 8	Г			
Nectarine			Γ	Г	•				0	0		0	•	•	0	•	0	9)		•	0	•	•	0	0	•	0	•	0	0	•	0	0		9	•	Г				Г	Γ			Γ	Г		Г	
Plum						10.									•		0	•		9		9	•	0	0	9	0	0	9	0	0	9	-	0			•													
Apple - Pink Lady			Γ	Γ	Г	Γ					T							0			9	0	•	•	0	•		0	•	0	0	•		0	0	0	0	•		0	•	0	•				Γ		Г	
Feijoa Quince	35						0010										100		Ì						0			0	•	0	•	0	0	0	0		0	0	9	0				8						
Grapefruit - Ruby	•	•	•	•	0					0	•	0	•		0			Γ		I										П	0	•	•	•		•	•	•		•	•	0	0	•	0		•	0		
Mandarin - Imperial	0	0	0	0	•	•	0		Ì																														9				6	6	0	0	8	0		9
Navel - Leng	•																																				0	•	•	0		0	0		0	0	•	0		

NOTE: Ripening periods will vary according to fruit variety, location and season.

PRE-HARVEST

- · Install monitoring traps & monitor weekly;
- · If QFF are found in traps apply weekly bait sprays; or
- Install insect nets / paper bags (the earlier the better).

RIPENING AND HARVEST

- · Monitor traps and check fruit for sting marks;
- · Continue weekly bait sprays;
- · Repair holes in nets and keep the net off the fruit;
- Destroy fallen and infested fruit (plastic bag / slash / deep burial);
- Pick fruit as it ripens and don't let it over-ripen on the tree;
- · Replace/recharge monitoring trap every 3 months.

POST-HARVEST

- · Continue weekly bait sprays for 2-3 weeks;
- · Remove and repair nets and move to another tree;
- Strip pick all remaining fruit and destroy (plastic bag / slash / deep burial)

